

RESEARCH ARTICLE

View Article Online
View Journal | View IssueCite this: *Org. Chem. Front.*, 2022, 9, 6490

"Thiol-free synthesized" and sustainable thiolating synthons for nickel-catalyzed reductive assembly of sulfides with high efficiency†

Yong Liu,^a Yuenian Xu,^a Yan Zhang,^a Wen-Chao Gao ^b and Xinxin Shao ^{*a}

Unsymmetrical sulfides are widely found in the pharmaceutical industry, organic synthesis, and materials science. As a result, it will be of great significance to discover mild and efficient methodologies and electrophilic sulfur transfer reagents, avoiding the general employment of odorous and toxic thiols. Herein, we present a highly efficient nickel-catalyzed cross-electrophile coupling of organic halides with "thiol-free" synthesized *N*-thiophthalimides as direct thiolating surrogates. This practical strategy features extremely low catalyst loading, good functional group tolerance and diverse downstream synthesis, enabling the construction of a broad range of sulfides under "base-free" conditions. Notably, a modified and more efficient reductive thiolation of disulfides was developed by employing our mild reaction conditions.

Received 17th August 2022,
Accepted 30th September 2022

DOI: 10.1039/d2qo01317g

rsc.li/frontiers-organic

Introduction


Sulfides, especially unsymmetrical sulfides, are central scaffolds that have found numerous applications in biological chemistry,¹ organic synthesis,² and materials science.³ As a consequence, it will be of great value to forge C–S bonds under mild reaction conditions. Classic nucleophilic substitutions of organic halides by thiols⁴ and advanced methods *via* radical pathways promoted by photo⁵ or electro-chemistry⁶ have already been achieved during the last few decades. However, odorous and toxic thiols, strong bases and/or harsh reaction conditions such as high temperature were the major limitations in the above strategies (Scheme 1a). To overcome the drawbacks of the limited thiols, several powerful electrophilic sulphur transfer reagents have been well developed.⁷ However, a majority of them were indeed prepared from the corresponding thiols or disulfides, and step-economy was sacrificed (Scheme 1b). Organolithium and organomagnesium reagents are air and moisture sensitive though they have been well utilized in the transition-metal-catalyzed cross-couplings with electrophilic sulphur synthons.^{7a–d}

Nevertheless, electrophilic sulphur reagents could participate in metal-catalyzed cross-electrophile couplings⁸ with


various carbon atom-centred electrophiles (Scheme 1b).⁹ From this viewpoint, thiol-free synthesized cationic thiolating reagents instead of toxic thiols will be a promising alternative in the construction of C–S bonds. Very recently, Wang and co-workers disclosed that shelf-stable benzimidazolium sulfonamides (IMDN-SO₂R) which were prepared from readily accessible sulfonyl chlorides could act as novel cationic thiolating precursors for nickel-catalyzed cross-electrophile coupling with various organic halides (Scheme 1b, left).^{9d}

Selective formation of cross-coupled products has been the long-standing issue existing in cross-electrophile coupling

a) Strategies for forging a C–S bond from different precursors.



b) Sulphur transfer reagents utilized in Cross-electrophile couplings.



Scheme 1 Selective strategies and thiolating reagents for C–S bond formation.

^aCollege of Material, Chemistry and Chemical Engineering, Key Laboratory of Organosilicon Chemistry and Material Technology of Ministry of Education, Hangzhou Normal University, Hangzhou 311121, Zhejiang, People's Republic of China. E-mail: xxshao@hznu.edu.cn

^bCollege of Biomedical Engineering, Taiyuan University of Technology, Taiyuan 030024, Shanxi, People's Republic of China

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d2qo01317g>


because of the competition of oxidative addition and transmetalation between the two coupling partners.^{8a,b} As a result, the disulfides are generated more easily as the major side products and indeed, homo-coupling of the electrophilic sulphur reagents is always observed in many cases. Moreover, a long reaction time and/or elevated reaction temperature were required to achieve good conversions and the scope was limited (Scheme 1b, left).^{9b,c}

Based on the significant achievements in the reductive cross-coupling for assembling C–C/C–X (X = Si, Ge, S, Se) bonds¹⁰ with our continuing interest¹¹ in the development of efficient reagents in synthetic chemistry, we speculate that *N*-thiophthalimides can serve as ideal precursors due to the weak “N–S” bond and easy modification of the reactivity by replacement of substituents on the reagent’s skeleton (Scheme 1c). *N*-Thiophthalimides are also air and moisture stable solids and can be further subjected to cross-electrophilic coupling. Thus, this work will provide direct information on “N–S⁺” type reagents in the reductive thiolation, while the activated “N–S⁺” intermediate was only confirmed by HRMS in the previous study.^{9d} Thus, a facile and “thiol-free” method¹² for preparation of such powerful reagents will directly enable the synthesis of functionalized sulfides under non-basic reaction conditions. More importantly, a sustainable strategy by recycling phthalimide as the original starting material after simple work-up will be developed, providing a good alternative as the electrophile in the C–S bond formation. Herein, we report our efforts on the synthesis of diversified *N*-thiophthalimides especially alkyl substituted ones from organozinc reagents¹³ followed by transformation towards the reductive cross-coupling with a variety of organic halides.

Results and discussion

Inspired by the pioneering work of Cornella’s group on the synthesis of *N*-arylthiophthalimides,¹⁴ we first tested thiol-free synthesis from easily accessible starting materials. Simply refluxing di(1-phthalimidyl)disulfane **1** in SO₂Cl₂ gave *N*-(chlorosulfonyl)phthalimide **2** in quantitative yield, while continuous bubbling of Cl₂ gas for a long time¹⁵ or an additive such as pyridine¹⁶ were required in previous studies (Table 1). On treatment with freshly-made organozinc reagents for a short time, a number of *N*-arylthiophthalimides containing hydrogen (**4a**), OMe (**4b**), Me (**4c**), F (**4d**), Br (**4e**), and Cl (**4f**) were formed in moderate to high yields after simple silica gel chromatography purification (Table 1). Notably, alkyl organozinc reagents easily generated from the corresponding alkyl bromides in the presence of zinc could react with compound **2** as well, generating challenging *N*-alkylthiophthalimides (**4g–i**) which were rarely explored¹⁴ in moderate yields. Finally, these *N*-thiophthalimides were isolated as shelf-stable solids and could be further stored in a refrigerator for several months without detectable decomposition according to ¹H NMR spectroscopy.

Table 1 Thiol-free synthesis of *N*-thiophthalimides^a



^a Reaction conditions: **2** (1.67 mmol), organozinc reagent **3a–f** (1.0 equiv.), CH₂Cl₂ (2.5 mL) at 0 °C for 20 min, and then at 22 °C for 0.5 h. ^b Reaction conditions: **2** (1.2 equiv.), organozinc reagent **3g–i** (3.3 mmol), CH₂Cl₂ (2.5 mL) at 0 °C for 20 min, and then at 22 °C for 2.0 h. Isolated yields.

With these readily available reagents in hand, we then investigated the nickel catalyzed reductive thiolation of aryl (hetero) halides with *N*-thiophthalimides we synthesized since the nickel catalyst exhibited a unique ability in such kind of transformations. *N*-Phenylthiophthalimide **4a** and 4-iodoanisole **5a** were chosen as the model substrates for the reaction optimization. Fortunately, the transformation proceeded rather well in the presence of NiCl₂(PPh₃)₂ as the catalyst, 1,10-phen **L1** as the ligand, and zinc as the reductant, furnishing thiolated product **6a** in >99% yield (Table 2, entry 1). As a result, it has been one of the most efficient protocols until now in the reductive thiolation. However, other transition-metal salts such as Cu(OTf)₂, CrCl₃ and PdCl₂(PPh₃)₂ were ineffective (Table 2, entries 2–4). Bipyridine type ligands showed better catalytic activity, enabling the formation of sulfide **6a** in high yield while other ligands such as PPh₃ failed to afford the product (Table 2, entries 5–7). To our delight, only 0.5 mol% NiCl₂(PPh)₂ and 1.0 mol% 1,10-phen exhibited excellent catalytic activity to give the sulfide **6a** in quantitative yield (Table 2, entry 8). Lower conversion of 4-iodoanisole **1a** was obtained when Mn was employed as the reductant or only 1.0 equiv. of zinc power were used (Table 2, entries 9 and 10). Moreover, no desired product was detected in the absence of zinc with full recovery of the starting materials (Table 2, entry 11). A slightly lower yield of **6a** was achieved when the reaction was conducted in a shorter time (Table 2, entry 12). Switching DMF to less polar solvents such as THF and CH₂Cl₂ led to low

Table 2 Optimization of the reaction conditions for the nickel-catalyzed reductive coupling of **4a** with **5a**^a

Entry	Catalyst	Ligand	Reductant	Solvent	x (equiv.)	T (°C)	Yield (%)
1	NiCl ₂ (PPh ₃) ₂	L1	Zn	DMF	2.0	50	>99
2	Cu(OTf) ₂	L1	Zn	DMF	2.0	50	<5
3	CrCl ₃	L1	Zn	DMF	2.0	50	<5
4	PdCl ₂ (PPh ₃) ₂	L1	Zn	DMF	2.0	50	<5
5	NiCl ₂ (PPh ₃) ₂	L2	Zn	DMF	2.0	50	88
6	NiCl ₂ (PPh ₃) ₂	L3	Zn	DMF	2.0	50	92
7	NiCl ₂ (PPh ₃) ₂	L4	Zn	DMF	2.0	50	<5
8	NiCl ₂ (PPh ₃) ₂	L1	Zn	DMF	2.0	22	>99
9	NiCl ₂ (PPh ₃) ₂	L1	Mn	DMF	2.0	22	<5
10 ^b	NiCl ₂ (PPh ₃) ₂	L1	Zn	DMF	2.0	22	<5
11	NiCl ₂ (PPh ₃) ₂	L1	—	DMF	1.2	22	<5
12 ^c	NiCl ₂ (PPh ₃) ₂	L1	Zn	DMF	1.2	22	98
13	NiCl ₂ (PPh ₃) ₂	L1	Zn	THF	1.2	22	<5
14	NiCl ₂ (PPh ₃) ₂	L1	Zn	CH ₂ Cl ₂	1.2	22	<5
15	NiCl ₂ (PPh ₃) ₂	L1	Zn	DMAc	1.2	22	>99
16 ^d	NiCl ₂ (PPh ₃) ₂	L1	Zn	DMF	1.2	22	<5

L1

L2

L3

L4

^a For entries 1–7: catalyst (2.5 mol%), ligand (3.0 mol%); for entries 8–16: NiCl₂(PPh₃)₂ (0.5 mol%), ligand L1 (1.0 mol%). ^b 1.0 equiv. of Zn was used. ^c 1 h. ^d Under air.

conversion, but DMAc, an analogue of DMF, gave the final sulfide **6a** in comparable yield (Table 2, entries 13–15). It was found that the reaction was totally inhibited in the presence of air (Table 2, entry 16).

Under identified conditions, the generality of the current methodology in the reductive thiolation was studied (Table 3). Pleasingly, a variety of functional groups including simple hydrogen (**6a**), methoxyl (**6b**), methyl (**6c**), fluoride (**6d**), bromide (**6e**), and chloride (**6f**) were well tolerated and unsymmetrical sulfides were obtained in moderate to high yields. Moreover, the reaction of alkyl *N*-thiophthalimides (**4g–i**) occurred as well when harsher reaction conditions were employed, affording the alkyl aryl sulfides (**6g–i**) in moderate to good yields.

Encouraged by the previously successful attempts, we next explored the versatility of carbon atom-centred electrophiles such as organic halides. It was found that the reaction of aryl (hetero) iodides proceeded smoothly in the presence of only 0.5 mol% NiCl₂(PPh₃)₂, 1.0 mol% 1,10-phen L1, and 2.5 equiv. of zinc, and the thiolated products were generated in reasonable yields after 2 h at 22 °C. Aryl iodides with functional groups such as methoxyl, chloride, bromide, and trifluoromethyl reacted to deliver the corresponding sulfides **6a**, **7a–b**, and **7i** in high yields (Table 4). Interestingly, substrates bearing the tosyl group **7c** and the boronate group **7d** at the *para*-position of the benzene ring were compatible with the reaction conditions, allowing the downstream synthesis of

Table 3 Scope of the nickel-catalyzed reductive coupling of **4b–i** with **5a**^a

<p>6b, 91% yield</p> <p>6d, 88% yield</p> <p>6f, 80% yield</p> <p>6h, 81% yield^b</p>	<p>6c, 89% yield</p> <p>6e, 81% yield</p> <p>6g, 98% yield^b</p> <p>6i, 96% yield^b</p>
---	---

^a Reaction conditions: **5a** (0.5 mmol), reagent **4a–f** (1.2 equiv.), NiCl₂(PPh₃)₂ (0.5 mol%), 1,10-phen (1.0 mol%), Zn (2.5 equiv.), and DMF (2.5 mL) at 22 °C for 2 h. Isolated yields are reported. ^b Reaction conditions: **5a** (0.5 mmol), reagent **4g–i** (2.0 equiv.), NiCl₂(PPh₃)₂ (5.0 mol%), 1,10-phen (10 mol%), Zn (3.0 equiv.), and DMF (0.5 mL) at 120 °C for 4 h. Isolated yields.

Table 4 Scope of the nickel-catalyzed reductive coupling of **4a** with **5a–w**^a




4a , 1.2 equiv	5a-w	6a, 7a-v
X = I		
6a , 87% yield	7a , 93% yield	7b , 74% yield
7c , 85% yield ^b	7d , 66% yield	7e , 97% yield
7f , 89% yield	7g , 81% yield	7h , 29% yield
7i , 83% yield (X = I)	7j , 79% yield (X = I)	7k , 93% yield (X = I)
7l , 82% yield (X = Br)	7m , 42% yield (X = Br)	7n , 81% yield (X = Br)
X = Br^d		
7o , 56% yield	7p , 84% yield	7q , 62% yield
7r , 50% yield	7s , 63% yield	7t , 80% yield
		6a , <5% yield
7u , 52% yield	7v , 46% yield	

^a Reaction conditions: **5a–w** (0.5 mmol), reagent **4a** (1.2 equiv.), NiCl₂(PPh₃)₂ (0.5 mol%), 1,10-phen (1.0 mol%), Zn (2.5 equiv.), and DMF (2.5 mL) at 22 °C for 2 h. ^b NiBr₂ (2.5 mol%), 1,10-phen (5.0 mol%), NaI (2.0 equiv.), Zn (3.0 equiv.), and DMF (2.5 mL) at 80 °C for 8 h. Isolated yields.

functional derivatives. Heteroaryl iodides including benzofuran **7e** and indole **7f** were engaged in the reductive cross-coupling with electrophilic thiolating reagent **4a** in high yields. Aryl iodides bearing carbonyl groups such as aldehyde **7j** and ester **7k** which were sensitive toward nucleophilic organolithium and organomagnesium reagents reacted readily to lead to the construction of C–S bonds efficiently. It was discovered that alkenyl and alkynyl iodides **7g–h** could readily convert into the thiolated products, resulting in the formation of the coupled products in acceptable yields. Notably, similar results were also achieved when aryl(hetero) bromides were applied as the coupling partners under the slightly modified reaction conditions. Reactions of aryl bromides containing the ester installed at the different position of the benzene ring as

well as strained cyclopropane substituted one led to the coupled products **7l–r** in reasonable yields. Moreover, a series of hetero(aryl) bromides could be successfully transformed into the desired sulfides **7s–v** in moderate to good yields, illustrating the applicability of this mild methodology. Unfortunately, the reaction of 4-chloroanisole **5w** failed to give any coupled product and further optimization of the reaction conditions was needed.

To further show the potential applications of the current protocol, late-stage thiolation of biologically complicated drug molecules and downstream functionalizations were conducted. Drug molecules fenofibrate and vitamin E derived aryl iodides could be successfully incorporated to afford the thiolated analogues **8** and **9** in high yields (Scheme 2a). A scale-up reaction proceeded smoothly, furnishing the coupled sulfides **7d** and

a) Late-stage modification of structurally complicated molecules.**b)** Scale-up synthesis of compounds **7d** and **7w**.**c)** Further derivations of the corresponding products (left: from **7d**, right: from **7w**).


Scheme 2 Late-stage modification of complex molecules and synthetic applications. ^a **5x–y** (0.5 mmol) or **5d** and **5z** (0.3 mmol) were used under the optimized conditions for **6a–7d** (0.20 mmol), AgNO₃ (0.060 equiv.), Et₃N (1.0 equiv.), EtOH/H₂O at 80 °C for 1 h in air. ^b **7d** (0.20 mmol), KHF₂ (5.0 equiv.), MeCN/H₂O at 30 °C for 9 h. ^c **7d** (1.2 equiv.), PhI (0.20 mmol), Pd(PPh₃)₄ (10 mol%), K₂CO₃ (2.0 equiv.), dioxane at 100 °C for 12 h. ^d **7w** (0.50 mmol), alkyne (1.2 equiv.), PdCl₂(PPh₃)₂ (5.0 mol%), K₂CO₃ (2.0 equiv.), Et₃N/DMF at 100 °C for 12 h. ^e **7w** (0.50 mmol), NiCl₂(PPh₃)₂ (10 mol%), 5,5'-dimethyl-2,2'-bipyridine (20 mol%), Mn (2.5 equiv.), chlorosilane (2.0 equiv.), DMF at 22 °C for 24 h. ^f **7w** (0.50 mmol), PhI(OAc)₂ (2.5 equiv.), (NH₄)₂CO₃ (2.0 equiv.), MeOH at 22 °C for 12 h. Isolated yields.

7w with excellent efficiency (Scheme 2b). Further transformations of the products were also explored since both of them were useful building blocks in organic synthesis (Scheme 2c). Thus, the boronic acid ester (R-Bpin) **7d** could readily convert into more stable trifluoroborate potassium salts (R-BF₃K) **11** *via* cleavage of the C–B bond.¹⁷ Suzuki coupling as well as protonation occurred to deliver the corresponding product **12** in moderate to good yields. Likewise, the derivative **7w** could react with the terminal alkyne *via* classic Sonogashira coupling, generating the corresponding product in 90% yield. Based on Shu's pioneering work,¹⁸ compound **7w** underwent a practical cross-electrophile coupling with vinyl chlorosilanes to forge a C(sp²)-Si bond. Moreover, the same compound could convert into sulfoximine **15** in high yield by treatment of (diacetoxyiodo)benzene and ammonium carbonate.^{11a}


This mild method also features sustainable utilization of the original starting material phthalimide **16**, which was isolated in 99% yield with >99% recovery by filtration after evaporation of the extracted solvents (Scheme 3a). Meanwhile, the thiolated product **6a** was obtained directly in 99% yield with >99% purity from the crude mixture after simple work-up on a 2.0 mmol scale (for more details, see the ESI†). *N*-Phenylthiophthalimide **4a**, the coupling partner, could be fully synthesized in three steps from the recycled phthalimide (Scheme 3b). Therefore, it provided a sustainable and practical strategy to construct a series of unsymmetrical sulfides under mild reaction conditions.

To better understand the reaction mechanism of this method, we then carried out the following experiments includ-

ing radical trapping, a control reaction and a stoichiometric reaction of the ArNi(II)I complex. First of all, no desired product **6a** was detected with >90% recovery of **5a** when TMEPO, a general radical scavenger, was added to the mixture (Scheme 4a). While in the cases of BHT and 9,10-dihydroanthracene, the desired product **5a** could still be obtained in 95% and >99% ¹H NMR yield, respectively, indicating that a radical pathway might not be involved. To probe whether disulfides could serve as thiolating surrogates in the reaction, we applied several disulfides to the process under the same reaction conditions employing *N*-thiophthalimides. Surprisingly, the transformation occurred smoothly to afford the corresponding sulfides **18a–d** under much milder reaction conditions (compared to Taniguchi's conditions) with higher efficiency (Scheme 4b). Thus, the disulfides were proved to be quite reactive in the reductive thiolation. Meanwhile, the



Scheme 3 Scale-up reaction and recycling of phthalimide **16**.



Scheme 4 Mechanistic investigations.




Fig. 1 Plausible mechanism of the current nickel-catalyzed reductive thiolation employing *N*-thiophthalimides.

corresponding disulfide **17c** was obtained in 40% NMR yield with full conversion of *N*-phenylthiophthalimide **4a** in a short time (10 min) when the reaction was conducted in the absence of aryl halides at 0 °C (Scheme 4c). A stoichiometric reaction of the $Ar-Ni(II)-I$ complex **18** with *N*-phenylthiophthalimide **4a** was then examined in the presence or absence of Zn (Scheme 4d) and the thiolated product **20** was generated in 57% 1H NMR yield.

Based on the above experimental results and previous reports,^{9d,19} a plausible catalytic cycle for this current nickel-catalyzed reductive thiolation is proposed as illustrated in Fig. 1. Initially, “*in situ*” formed $L_nNi(0)$ **I** undergoes oxidative addition of the C–X bond of aryl halides (R–X) to form the intermediate **II**, and after reduction by Zn, the $L_nNi(I)R$ **III** was generated. After treatment of disulfides which were formed from the corresponding *N*-thiophthalimides, the $L_nNi(I)R$ **III** can convert into intermediate **IV** before the formation of the adduct **IV**. Finally, reductive elimination of $L_nNi(III)R$ **III** occurs to afford the final product and $L_nNi(I)R$ **VI**. Upon reduction with Zn(0), catalytically active $L_nNi(0)$ **I** is regenerated and facilitates the next catalytic cycle.

Conclusions

In summary, we successfully developed an efficient, “thiol-free” and sustainable strategy to prepare a series of *N*-thiophthalimides from aryl and alkylzinc reagents. A nickel-catalyzed reductive thiolation of organic halides with bench-stable and readily accessible *N*-thiophthalimides as powerful electrophilic sulphur reagents was also documented. This non-basic process can proceed well under extremely mild reactions promoted by as little as 5.0 mol% loading of nickel catalyst and is further highlighted by late-stage modification of biologically complicated drug molecules, downstream synthesis of related important compounds and sustainable use of the

original starting materials. Moreover, direct evidence of the high reactivity of the “N–S” type sulphur transfer reagents in the reductive formation of C–S bonds was explored. Thus, our strategy provided a practical, straightforward and modular platform to assemble a variety of unsymmetrical sulfides in good to excellent yields. Investigation of more challenging substrates such as alkyl halides or unactivated alcohols is under way and the results will be reported in the near future.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We thank the National Science Foundation of China (22001051), the Hangzhou Leading Innovation and Entrepreneurship Team Project (TD2020015) and the Start-up Funding of Hangzhou Normal University for financial support. We also greatly thank Dr Chunqi Sheng (Hangzhou Normal University) and Dr Hong Dong (Hangzhou Normal University) for the assistance with IR and HRMS analyses. We thank Prof Pengfei Zhang for assistance and helpful discussions.

References

- (a) K. L. Dunbar, D. H. Scharf, A. Litomska and C. Hertweck, Enzymatic Carbon-Sulfur Bond Formation in Natural Product Biosynthesis, *Chem. Rev.*, 2017, **117**, 5521–5577; (b) N. Wang, P. Saidhareddy and X. Jiang, Construction of Sulfur-containing Moieties in The Total Synthesis of Natural Products, *Nat. Prod. Rep.*, 2020, **37**, 246–275; (c) M. Feng, B. Tang, S. Liang and X. Jiang, Sulfur Containing Scaffolds in Drugs: Synthesis and Application in Medicinal Chemistry, *Curr. Top. Med. Chem.*, 2016, **16**, 1200–1216.
- (a) M. Wang, C. Wang and X. Jiang, Recent Progress in the Sulfur-Containing Perfume & Flavors Construction, *Chin. J. Org. Chem.*, 2019, **39**, 2139–2147; (b) V. K. Aggarwal and C. L. Winn, Catalytic, Asymmetric Sulfur Ylide-Mediated Epoxidation of Carbonyl Compounds: Scope, Selectivity, and Applications in Synthesis, *Acc. Chem. Res.*, 2004, **37**, 611–620; (c) M. Mellah, A. Voituriez and E. Schulz, Chiral Sulfur Ligands for Asymmetric Catalysis, *Chem. Rev.*, 2007, **107**, 5133–5209.
- (a) D. A. Boyd, Sulfur and Its Role In Modern Materials Science, *Angew. Chem., Int. Ed.*, 2016, **55**, 15486–15502; (b) K. Takimiya, S. Shinamura, I. Osaka and E. Miyazaki, Thienoacene-Based Organic Semiconductors, *Adv. Mater.*, 2011, **23**, 4347–4370.
- (a) E. Alvaro and J. F. Hartwig, Resting State and Elementary Steps of the Coupling of Aryl Halides with Thiols Catalyzed by Alkylbisphosphine Complexes of Palladium, *J. Am. Chem. Soc.*, 2009, **131**, 7858–7868; (b) P. Bichler and J. A. Love, in *C–X Bond Formation*, ed. A.

- Vigalok, Springer Berlin Heidelberg, Berlin, Heidelberg, 2010, pp. 39–64, DOI: [10.1007/978-3-642-12073-2_3](https://doi.org/10.1007/978-3-642-12073-2_3); (c) C.-F. Lee, Y.-C. Liu and S. S. Badsara, Transition-Metal-Catalyzed C-S Bond Coupling Reaction, *Chem. – Asian J.*, 2014, **9**, 706–722; (d) B. Xiong, X. Zeng, S. Geng, S. Chen, Y. He and Z. Feng, Thiyl radical promoted chemo- and regioselective oxidation of C=C bonds using molecular oxygen via iron catalysis, *Green Chem.*, 2018, **20**, 4521–4527; (e) J. You, Q. Chen and Y. Nishihara, Nickel-Catalyzed Decarbonylative Thioetherification of Acyl Fluorides via C-F Bond Activation, *Synthesis*, 2021, **53**, 3045–3050.
- 5 B. Liu, C.-H. Lim and G. M. Miyake, Visible-Light-Promoted C-S Cross-Coupling via Intermolecular Charge Transfer, *J. Am. Chem. Soc.*, 2017, **139**, 13616–13619.
- 6 (a) D. Liu, H.-X. Ma, P. Fang and T.-S. Mei, Nickel-Catalyzed Thiolation of Aryl Halides and Heteroaryl Halides through Electrochemistry, *Angew. Chem., Int. Ed.*, 2019, **58**, 5033–5037; (b) F. Zhang, Y. Wang, Y. Wang and Y. Pan, Electrochemical Deoxygenative Thiolation of Preactivated Alcohols and Ketones, *Org. Lett.*, 2021, **23**, 7524–7528; (c) Y. Wang, L. Deng, X. Wang, Z. Wu, Y. Wang and Y. Pan, Electrochemically Promoted Nickel-Catalyzed Carbon-Sulfur Bond Formation, *ACS Catal.*, 2019, **9**, 1630–1634.
- 7 (a) T. Kondo and T.-a. Mitsudo, Metal-Catalyzed Carbon-Sulfur Bond Formation, *Chem. Rev.*, 2000, **100**, 3205–3220; (b) C. Savarin, J. Srogl and L. S. Liebeskind, A Mild, Nonbasic Synthesis of Thioethers. The Copper-Catalyzed Coupling of Boronic Acids with N-Thio(alkyl, aryl, heteroaryl)imides, *Org. Lett.*, 2002, **4**, 4309–4312; (c) B.-X. Du, Z.-J. Quan, Y.-X. Da, Z. Zhang and X.-C. Wang, Chemo-Controlled Cross-Coupling of Di(hetero)aryl Disulfides with Grignard Reagents: C-C vs. C-S Bond Formation, *Adv. Synth. Catal.*, 2015, **357**, 1270–1276; (d) S. Chen, Q. Wen, Y. Zhu, Y. Ji, Y. Pu, Z. Liu, Y. He and Z. Feng, Boron-promoted Reductive Deoxygenation Coupling Reaction of Sulfonyl Chlorides for the C(sp³)-S Bond Construction, *Chin. Chem. Lett.*, 2022, **33**, 5101–5105; (e) M. Iwasaki, W. Kaneshika, Y. Tsuchiya, K. Nakajima and Y. Nishihara, Palladium-Catalyzed peri-Selective Chalcogenation of Naphthylamines with Diaryl Disulfides and Diselenides via C-H Bond Cleavage, *J. Org. Chem.*, 2014, **79**, 11330–11338; (f) M. Iwasaki, M. Iyanaga, Y. Tsuchiya, Y. Nishimura, W. Li, Z. Li and Y. Nishihara, Palladium-Catalyzed Direct Thiolation of Aryl C-H Bonds with Disulfides, *Chem. – Eur. J.*, 2014, **20**, 2459–2462.
- 8 (a) D. J. Weix, Methods and Mechanisms for Cross-Electrophile Coupling of Csp² Halides with Alkyl Electrophiles, *Acc. Chem. Res.*, 2015, **48**, 1767–1775; (b) D. A. Everson and D. J. Weix, Cross-Electrophile Coupling: Principles of Reactivity and Selectivity, *J. Org. Chem.*, 2014, **79**, 4793–4798; (c) J. Gu, X. Wang, W. Xue and H. Gong, Nickel-catalyzed Reductive Coupling of Alkyl Halides with Other Electrophiles: Concept and Mechanistic Considerations, *Org. Chem. Front.*, 2015, **2**, 1411–1421; (d) S. Li, Y. Wang, L. Zhong, S. Wang, Z. Liu, Y. Dai, Y. He and Z. Feng, Boron-Promoted Umpolung Reaction of Sulfonyl Chlorides for the Stereospecific Synthesis of Thioglycosides via Reductive Deoxygenation Coupling Reactions, *Org. Lett.*, 2022, **24**, 2463–2468; (e) Y. Feng, S. Yang, S. Zhao, D.-P. Zhang, X. Li, H. Liu, Y. Dong and F.-G. Sun, Nickel-Catalyzed Reductive Aryl Thiocarbonylation of Alkene via Thioester Group Transfer Strategy, *Org. Lett.*, 2020, **22**, 6734–6738.
- 9 (a) N. Taniguchi and T. Onami, Magnesium-Induced Copper-Catalyzed Synthesis of Unsymmetrical Diaryl Chalcogenide Compounds from Aryl Iodide via Cleavage of the Se-Se or S-S Bond, *J. Org. Chem.*, 2004, **69**, 915–920; (b) N. Taniguchi, Alkyl- or Arylthiolation of Aryl Iodide via Cleavage of the S-S Bond of Disulfide Compound by Nickel Catalyst and Zinc, *J. Org. Chem.*, 2004, **69**, 6904–6906; (c) P. Mao, H.-D. Sui, J.-W. Yuan, Y.-M. Xiao, L.-R. Yang and W.-P. Mai, Nickel-Catalyzed Carbon-Sulfur Bond Formation through Couplings of Aryl Iodides and Aryl Ethanethioates, *ChemistrySelect*, 2020, **5**, 9908–9910; (d) W. Zhang, M. Huang, Z. Zou, Z. Wu, S. Ni, L. Kong, Y. Zheng, Y. Wang and Y. Pan, Redox-active Benzimidazolium Sulfonamides as Cationic Thioliating Reagents for Reductive Cross-coupling of Organic Halides, *Chem. Sci.*, 2021, **12**, 2509–2514; (e) S. Yoshida, Y. Sugimura, Y. Hazama, Y. Nishiyama, T. Yano, S. Shimizu and T. Hosoya, A Mild and Facile Synthesis of Aryl and Alkenyl Sulfides via Copper-catalyzed Deborylthiolation of Organoborons with Thiosulfonates, *Chem. Commun.*, 2015, **51**, 16613–16616; (f) K. Kanemoto, Y. Sugimura, S. Shimizu, S. Yoshida and T. Hosoya, Rhodium-catalyzed Odorless Synthesis of Diaryl Sulfides from Borylarenes and S-aryl thiosulfonates, *Chem. Commun.*, 2017, **53**, 10640–10643; (g) P. Mampuy, Y. Zhu, T. Vlaar, E. Ruijter, R. V. A. Orru and B. U. W. Maes, Sustainable Three-Component Synthesis of Isothioureas from Isocyanides, Thiosulfonates, and Amines, *Angew. Chem., Int. Ed.*, 2014, **53**, 12849–12854; (h) Y. Dong, P. Ji, Y. Zhang, C. Wang, X. Meng and W. Wang, Organophotoredox-Catalyzed Formation of Alkyl-Aryl and -Alkyl C-S/Se Bonds from Coupling of Redox-Active Esters with Thio/Selenosulfonates, *Org. Lett.*, 2020, **22**, 9562–9567.
- 10 (a) X. Pang, P.-F. Su and X.-Z. Shu, Reductive Cross-Coupling of Unreactive Electrophiles, *Acc. Chem. Res.*, 2022, **55**(17), 2491–2509; (b) Z.-Z. Zhao, X. Pang, X.-X. Wei, X.-Y. Liu and X.-Z. Shu, Nickel-Catalyzed Reductive C(sp²)-Si Coupling of Chlorohydrosilanes via Si-Cl Cleavage, *Angew. Chem., Int. Ed.*, 2022, **61**, e202200215; (c) P.-F. Su, K. Wang, X. Peng, X. Pang, P. Guo and X.-Z. Shu, Nickel-Catalyzed Reductive C-Ge Coupling of Aryl/Alkenyl Electrophiles with Chlorogermanes, *Angew. Chem., Int. Ed.*, 2021, **60**, 26571–26576; (d) L. Qi, X. Pang, K. Yin, Q.-Q. Pan, X.-X. Wei and X.-Z. Shu, Mn-mediated Reductive C(sp³)-Si Coupling of Activated Secondary Alkyl Bromides with Chlorosilanes, *Chin. Chem. Lett.*, 2022, **33**, 5061–5064.
- 11 (a) Y. Liu, S. Xing, J. Zhang, W. Liu, Y. Xu, Y. Zhang, K. Yang, L. Yang, K. Jiang and X. Shao, Construction of Diverse C-S/C-Se Bonds via Nickel Catalyzed Reductive Coupling Employing Thiosulfonates and a Selenosulfonate

- under Mild Conditions, *Org. Chem. Front.*, 2022, **9**, 1375–1382; (b) S. Xing, Y.-Y. Zhu, W. Liu, Y. Liu, J. Zhang, H. Zhang, Y. Wang, S.-F. Ni and X. Shao, C–H Fluoroalkylsulfonylation/Intramolecular Rearrangement for Precise Synthesis of Fluoroalkyl Sulfoxides, *Org. Lett.*, 2022, **24**, 3378–3383.
- 12 (a) C. Xu and Q. Shen, Palladium-Catalyzed Trifluoromethylthiolation of Aryl C–H Bonds, *Org. Lett.*, 2014, **16**, 2046–2049; (b) F. Shen, P. Zhang, L. Lu and Q. Shen, [[[Ethoxycarbonyl]difluoromethyl]thio]phthalimide: A Shelf-Stable, Electrophilic Reagent with a Convertible Group for the Synthesis of Diversified Fluoroalkylthiolated Compounds, *Org. Lett.*, 2017, **19**, 1032–1035.
- 13 (a) P. Knochel and R. D. Singer, Preparation and Reactions of Polyfunctional Organozinc Reagents in Organic Synthesis, *Chem. Rev.*, 1993, **93**, 2117–2188; (b) J. Li, E. Tan, N. Keller, Y.-H. Chen, P. M. Zehetmaier, A. C. Jakowetz, T. Bein and P. Knochel, Cobalt-Catalyzed Electrophilic Aminations with Anthranils: An Expedient Route to Condensed Quinolines, *J. Am. Chem. Soc.*, 2019, **141**, 98–103; (c) S. Graßl, C. Hamze, T. J. Koller and P. Knochel, Copper-Catalyzed Electrophilic Thiolation of Organozinc Halides by Using N-Thiophthalimides Leading to Polyfunctional Thioethers, *Chem. – Eur. J.*, 2019, **25**, 3752–3755; (d) P. Saravanan and P. Anbarasan, Palladium Catalyzed Aryl(alkyl)thiolation of Unactivated Arenes, *Org. Lett.*, 2014, **16**, 848–851.
- 14 L. Wang and J. Cornella, A Unified Strategy for Arylsulfur (vi) Fluorides from Aryl Halides: Access to Ar-SOF₃ Compounds, *Angew. Chem., Int. Ed.*, 2020, **59**, 23510–23515.
- 15 (a) S. A. Hutchinson, S. P. Baker, J. Linden and P. J. Scammells, New Potent and Selective A1 Adenosine Receptor Agonists, *Biorg. Med. Chem.*, 2004, **12**, 4877–4884; (b) M. U. Bombala and S. V. Ley, Preparation of Episulphides from Alkenes via Succinimide-N-sulphenyl Chloride or Phthalimide-N-sulphenyl Chloride Adducts, *J. Chem. Soc., Perkin Trans. 1*, 1979, 3013–3016, DOI: [10.1039/P19790003013](https://doi.org/10.1039/P19790003013).
- 16 (a) D. Zhu, Y. Gu, L. Lu and Q. Shen, N-Difluoromethylthiophthalimide: A Shelf-Stable, Electrophilic Reagent for Difluoromethylthiolation, *J. Am. Chem. Soc.*, 2015, **137**, 10547–10553; (b) F. Xiong, L. Lu, T.-Y. Sun, Q. Wu, D. Yan, Y. Chen, X. Zhang, W. Wei, Y. Lu, W.-Y. Sun, J. J. Li and J. Zhao, A Bioinspired and Biocompatible *ortho*-Sulfiliminy Phenol Synthesis, *Nat. Commun.*, 2017, **8**, 15912.
- 17 G. A. Molander, Organotrifluoroborates: Another Branch of the Mighty Oak, *J. Org. Chem.*, 2015, **80**, 7837–7848.
- 18 J. Duan, K. Wang, G.-L. Xu, S. Kang, L. Qi, X.-Y. Liu and X.-Z. Shu, Cross-Electrophile C(sp²)-Si Coupling of Vinyl Chlorosilanes, *Angew. Chem., Int. Ed.*, 2020, **59**, 23083–23088.
- 19 (a) V. Gómez-Benítez, O. Baldovino-Pantaleón, C. Herrera-Álvarez, R. A. Toscano and D. Morales-Morales, High Yield Thiolation of Iodobenzene Catalyzed by the Phosphinite Nickel PCP Pincer Complex: [NiCl{C₆H₃-2,6-(OPPh₂)₂}], *Tetrahedron Lett.*, 2006, **47**, 5059–5062; (b) S. Biswas and D. J. Weix, Mechanism and Selectivity in Nickel-Catalyzed Cross-Electrophile Coupling of Aryl Halides with Alkyl Halides, *J. Am. Chem. Soc.*, 2013, **135**, 16192–16197.